

Four-Lined Plant Bug

Carla Staab*, Milwaukee Area Technical College, and Phil Pellitteri, UW Insect Diagnostic Lab

The adult four-lined plant bug (*Poecalocapsus lineatus*) is a 1/2 inch long, yellowish to yellowish-green true bug with four longitudinal black lines down the wing covers and black antennae. This plant bug looks somewhat like a striped cucumber beetle. Nymphs of the insect are wingless and bright yellow to red with rows of black spots on the abdomen. Older nymphs are yellowish-green with a yellow stripe on each wing pad.



An adult plant bug.



Four-lined plant bug damage.

Plants Attacked and Damage: Both adults and nymphs have piercing-sucking mouthparts and attack a wide variety of annuals, herbaceous perennials, woody shrubs, vegetables, and herbs including dogwood, forsythia, honeysuckle, hydrangea, viburnum, weigela, coreopsis, dahlia, mint, basil, morning glory, mums, lettuce and zinnia. On some hosts, feeding causes curling, distortion, or browning. On most plants feeding damage causes black or translucent, circular spots (about 1/16 inch in diameter) giving the plants a diseased appearance. Dead plant tissue may drop, leaving a 'shot hole' similar to some leaf spot diseases. When disturbed, plant bugs drop quickly to the ground and hide under foliage. Therefore, you may not actually see plant bugs causing plant damage. Damage is mainly cosmetic, but can be a concern on herbs and flowering plants.

Life Cycle: Four-lined plant bugs overwinter as eggs. Eggs hatch shortly after foliage appears in late April or early May, and nymphs begin feeding on the undersides of leaves. Nymphs feed for about thirty days and do most of the damage because they are relatively immobile compared to adults. Adults feed for about a month and then mate. Females cut slits lengthwise into the stems of woody or herbaceous plants and deposit one half dozen or more eggs into each slit. There is one generation per year.

Control: Begin monitoring plants by mid-May for signs of plant bug feeding. Hand picking can be effective if your eye and hand are faster than the plant bugs. Insecticidal soap can be used on edible plants and ornamentals. In addition, most garden insecticides are effective. In the fall, after the leaves drop, egg-containing slits can be easily seen on trees and shrubs, and these branches can be pruned out.

For more information on four-lined plant bug: Contact your county Extension agent.

*Completed as partial fulfillment of the requirements for an associate degree in Horticulture at the Milwaukee Area Technical College.

© 2002 by the Board of Regents of the University of Wisconsin System doing business as the division of Cooperative Extension of the University of Wisconsin Extension.

An EEO/Affirmative Action employer, University of Wisconsin Extension provides equal opportunities in employment and programming, including Title IX and ADA requirements. This document can be provided in an alternative format by calling Brian Hudelson at (608) 262-2863 (711 for Wisconsin Relay).

References to pesticide products in this publication are for your convenience and are not an endorsement or criticism of one product over similar products. You are responsible for using pesticides according to the manufacturer's current label directions. Follow directions exactly to protect the environment and people from pesticide exposure. Failure to do so violates the law.

Thanks to Karen Delahaut, Ann Joy, and Brian Hudelson for reviewing this document.

A complete inventory of University of Wisconsin Garden Facts is available at the University of Wisconsin-Extension Horticulture website: whort.uwex.edu.