2015 Winter Seminar Series

Top Ten Diseases of 2014

Brian D. Hudelson
Department of Plant Pathology
University of Wisconsin-Madison/Extension







2014 Winter Seminar Series Winter Injury

- Causes
 - Drought stress
 - Extreme winter conditions
- Affected plants
 - Virtually anything
 - Evergreens (yews and boxwoods)
 - Fruit trees
 - Redbud



2014 Winter Seminar Series Winter Injury

- Control
 - Water trees and shrubs adequately, particularly in the fall
 - Plant sensitive trees and shrubs in protected locations
 - Insulate sensitive plants where possible
 - Pray for snow


2014 Winter Seminar Series Verticillium Wilt

- Causes: <u>Verticillium dahliae</u> <u>Verticillium albo-atrum</u>
- Hosts
 - Many woody ornamentals
 - Common: Maple, ash, redbud, smokebush
 - "New": Seven son flower, wafer-ash, buttonbush
 - Many herbaceous plants
 - Many vegetables (tomato, potato, eggplant)
- Favorable environment: Cool, wet weather



2014 Winter Seminar Series Verticillium Wilt

- Control
 - Avoid Verticillium-infested areas
 - Pretest soils/mulches/composts for the presence of <u>Verticillium</u>
 - Fumigate heavily infested soils
 - Keep broad-leaf weeds under control
 - Avoid municipal mulches





• Wood Chips as an Inoculum Source

- Amur maple
- 30.0%/25.0% (Trted)
- 0.0%/0.0% (Non-Trted)
- Green Ash
- 23.7%/10.5% (Trted)
- 0.0%/0.0% (Non-Trted)
 - Redbud
- 10.7%/13.3% (Trted)
- 0.0%/0.0% (Non-Trted)

2014 Winter Seminar Series Verticillium Wilt


- Control
 - Use "resistant" plants
 - CONIFERS: Pines, spruces, firs, junipers
 - DECIDUOUS TREES/SHRUBS: Beech, birch, ginkgo, hackberry, hawthorn, hickory, honey locust, mountain ash, white oak, bur oak, poplar, serviceberry, sycamore, willow

2014 Winter Seminar Series Verticillium Wilt

- Control
 - Prevent plant stress
 - Prune diseased (wilted) areas
 - Decontaminate pruning tools
 - Make infected trees comfortable until they
 - Remove diseased plants
 - Destroy infected materials
 - Composting?

2014 Winter Seminar Series Guignardia Leaf Blotch

- Cause: Guignardia aesculi
- Hosts
 - Horse-chestnut
 - Ohio buckeye
- Favorable environment: Cool, wet weather



2014 Winter Seminar Series Guignardia Leaf Blotch


- Control
 - DO NOT panic
 - Remove diseased leaves
 - Use fungicides to prevent infections
 - Copper-containing fungicides, chlorothalonil, mancozeb, thiophanate methyl
 - 3 applications at bud break, 1/2 expansion of leaves, full leaf expansion

2014 Winter Seminar Series Tar Spot

• Causes: Rhytisma americanum Rhytisma acerinum

• Hosts: Maples

• Favorable environment: Cool, wet weather



2014 Winter Seminar Series Tar Spot

- Control
 - DO NOT panic
 - Remove diseased leaves
 - Burn
 - Bury
 - Hot compost
 - Use fungicides to prevent infections
 - Copper-containing fungicides
 - At bud break, 1/2 and full leaf expansion

2014 Winter Seminar Series Viral Diseases

- Causes
 - Many and varied
 - Tobacco mosaic virus (TMV)
 - Cucumber mosaic virus (CMV)
 - Impatiens necrotic spot virus (INSV)
 - Hosta virus X (HVX)
 - Cymbidium mosaic virus (CyMV)
 - Alfalfa mosaic virus (AMV)

2014 Winter Seminar Series Viral Diseases

- Hosts
 - Wide range: TMV, CMV, INSV, HVX, AMV
 - Narrow range: CyMV
- Favorable environment: None
- Transmission
 - Touch (TMV)
 - Mechanical injury (HVX, CyMV)
 - Insects (CMV, INSV, AMV)



2014 Winter Seminar Series Viral Diseases

- Control
 - Buy plants from a reputable source
 - DO NOT buy symptomatic plants
 - Pretest plants for viruses
 - Keep weeds under control
 - Control vectors (insects)
 - DO NOT smoke around your plants
 - Remove and destroy infected plants


2014 Winter Seminar Series Viral Diseases

- Control
 - Wash hands routinely
 - Disinfest contaminated materials
 - 1% sodium dodecyl sulfate (sodium lauryl sulfate) + 1% Alconox® (2½ Tbsp + 2¾ Tbsp/gal)
 - 20% low fat dry milk (Carnation®) + 0.1% polysorbate 20 (9% cups + ¾ tsp/gal)
 - Trisodium phosphate (14 dry oz/gal)
 - Alcohol dip followed by flaming

2014 Winter Seminar Series Impatiens Downy Mildew

- Cause: Plasmopara obducens
- Hosts
 - Standard garden impatiens (<u>I</u>. <u>walleriana</u>)
 - Balsam impatiens (<u>I. balsamina</u>)
 - Jewelweed (<u>I</u>. pallida, <u>I</u>. capensis)
 - New Guinea impatiens (<u>I</u>. <u>hawkeri</u>) (resistant/tolerant)
- Favorable environment: Wet weather





2014 Winter Seminar Series Impatiens Downy Mildew

- Control
 - Grow tolerant/resistant/immune plants
 - Start with clean transplants and seed
 - Keep materials from different sources physically separated
 - DO NOT grow impatiens in the same area every year
 - DO NOT overcrowd plants
 - DO NOT overhead water

2014 Winter Seminar Series Impatiens Downy Mildew

- Control
 - Watch for disease on a regular basis
 - Bag and discard affected plants
 - Symptomatic plants
 - Asymptomatic surrounding plants
 - Disinfest contaminated materials

2014 Winter Seminar Series Impatiens Downy Mildew

- Control
 - Use fungicides to prevent infections
 - Mefenoxam, fluopicolide, potassium phosphite, mancozeb, pyraclostrobin + boscalid, fluoxastobin, cyazofamid, dimethomorph, fenamidone, azoxystrobin
 - Alternate active ingredients (FRAC codes)
 - Apply at 7 day application intervals

2014 Winter Seminar Series Foliar Nematodes

- Cause: Aphelenchoides spp.
- Hosts
 - Wide host range
 - Hosta
 - Plants in the Saxifrage family
- Favorable environment: None





2014 Winter Seminar Series Foliar Nematodes

- Control
 - Start with clean propagation materials
 - Follow strict sanitation procedures when working with plant materials
 - Remove symptomatic plants
 - Remove contaminated plant debris
 - Avoid overhead irrigation
 - Hot water treatments (10 minutes at 125°F)

2014 Winter Seminar Series Southern Blight

- Pathogen: Sclerotium rolfsii
- Hosts
 - Many other herbaceous annuals and perennials
 - Some woody ornamentals
 - Hosta
 - Bedding plants
- Favorable environment: None



2014 Winter Seminar Series Southern Blight

- Control
 - Inspect plants prior to purchase and avoid infected plants
 - Avoid cocoa mulch (????)
 - Remove infected plants
 - Use fungicides for control
 - triadimefon
 - 14 28 day intervals
 - Pray for a cold winter

2014 Winter Seminar Series Plum Pockets

- Cause: <u>Taphrina communis</u>
- Host: Plum
- Favorable environment: Wet weather





2014 Winter Seminar Series Plum Pockets

- Control
 - Remove and destroy symptomatic fruits
 - Burn
 - Bury
 - Hot compost
 - Prune/thin trees to improve air flow
 - Use fungicides to prevent infections
 - Chlorothalonil, copper, ferbam
 - Apply after leaf fall and/or before leaf emergence

2014 Winter Seminar Series Rhizosphaera Needle Cast

- Pathogen: <u>Rhizosphaera kalkhoffii</u> (<u>Rhizosphaera sp.</u>)
- Hosts (major)
 - Colorado blue spruce
 - Other spruces: Engelmann, black, Serbian, Sitka

2014 Winter Seminar Series Rhizosphaera Needle Cast

- Hosts (minor)
 - Pines: Austrian, mugo, eastern white pine
 - Douglas fir
 - Hemlock
 - Balsam fir
- Favorable environment
 - Long periods of needle wetness
 - High humidity



2014 Winter Seminar Series Rhizosphaera Needle Cast

- Control
 - DO NOT plant Colorado blue spruce
 - DO NOT crowd trees when planting
 - Thin healthy branches to increase airflow
 - Prevent tree stress
 - Prune diseased branches

2014 Winter Seminar Series Rhizosphaera Needle Cast

- Control
 - Use fungicides to prevent infections
 - Copper-containing fungicides, chlorothalonil
 - Bud break
 - 3-4 week application interval under favorable conditions

2014 Winter Seminar Series How to Contact the PDDC

Plant Disease Diagnostics Clinic Department of Plant Pathology University of Wisconsin-Madison 1630 Linden Drive Madison, WI 53706-1598 (608) 262-2863 pddc@plantpath.wisc.edu http://pddc.wisc.edu Follow on Twitter @UWPDDC